

# Microlensing


Scott Gaudi (OSU), SOC



Andy Boden (MSC), LOC

J.P. Beaulieu (IAP), Dave Bennett (UND),  
Edward Cheng (Conceptual Analytics), Kem Cook  
(LLNL), Andy Gould (OSU), John Mather  
(NASA), Charley Noecker (Ball), Domenick  
Tenerelli (Lockheed)

# Microlensing Peculiarities

- Microlensing planet community is comparatively small.
- General consensus on forward directions.
- Two (and only two) paths forward
  - Ground based, 1-5 years, ~\$10-20M
    - Frequency of planets  $>M_{\oplus}$  beyond the snow line.
  - Space based, 5-10 years, ~\$300M
    - Complete census of planets with mass greater than Mars and  $a > 0.5$  AU, including habitable planets.






### Beyond the snow line:


- Location of giant planet formation (and our giant planets).
- ‘Failed Jupiters’
- Source of water

*Ground-based  $\mu$ lensing surveys probe planets with  $M > M_{\oplus}$  beyond the snow-line.*



*A space-based survey will provide a complete census of planetary systems with mass greater than Mars and  $a > 0.5$  AU (from 0 to  $\infty$  with Kepler), including habitable planets.*

# Detecting Planets



Primary event:


- Smooth, symmetric
- Typically 20 days

Planetary perturbation to images

- Short-timescale bump
- Measure:
  - Projected Separation
  - Mass Ratio

$$t_p = q^{1/2} t_E \approx 1 \text{ day} \left( \frac{M_p}{M_J} \right)^{1/2}$$


# Microlensing is directly sensitive to planet mass



- Works by perturbing images
- Does not require light from the lens or planet.
- Sensitive to planets in the disk and bulge with  $D_{OL}=1-8$  kpc
- Most sensitive to planets near the Einstein radius
- Sensitive to wide or free-floating planets
- Not sensitive to very close planets

# Very Low Mass Planets

Signal magnitude is *independent* of planet mass.




- Magnitude depends on separation of planet from image.
- Duration depends on mass.

$$t_p = q^{1/2} t_E \approx 2 \text{ hrs} \left( \frac{M_p}{M_{\odot}} \right)^{1/2}$$

- Signals get rarer and briefer.
- Detection Probability  
~ few %
- **Large** (~10%) signals for low-mass (Earth-mass) planets



# How Low Can We Go?



(Bennett & Rhie 1996)

- Limited by Source Size

$$\rho_* = \frac{\theta_*}{\theta_E} \approx 1$$


$$\theta_E \approx \mu\text{as} \left( \frac{M_p}{M_{\odot}} \right)^{1/2}$$



$$\theta_* \approx \mu\text{as} \left( \frac{R_*}{R_{\odot}} \right)$$

**Mars-mass planets detectable  
if solar-type sources can be  
monitored!**

# Sensitivity Depends Weakly on Host Mass



Sensitive to planets around:

- Main-sequence stars with  $M < M_\odot$
- Brown dwarfs
- Remnants


# Microlensing Strengths

- Peak sensitivity beyond the snow line.
  - 50-100K
- Sensitivity down to *very* low-mass planets.
  - Mass greater than that of ~10% Mars.
- Sensitivity to long-period and free-floating planets.
  - 0.5 AU -  $\infty$
- Sensitivity to planets over a wide range of host masses.
  - $M < M_{\odot}$
- Sensitivity to planets throughout the Galaxy.
  - 1-8 kpc
- Sensitivity to multiple-planet systems.

# Commonly heard complaints...

- But you don't know anything about the star, orbits, etc!
  - Typically can measure host star and planet masses to  $\sim 10\text{-}20\%$ .
  - In some special cases can learn something about the orbit.
- But the systems are so far away and faint!
  - Sufficiently bright to measure flux, color, and in some cases get spectra.
- But you only see it once!
  - Signals are large and unambiguous.
  - Demographics of planetary systems.


OGLE-2005-BLG-071  
(Udalski et al 2005)



$$M_p \sim 3.5 M_J, \quad r \sim 3.6 \text{ AU}$$

$$M_* \sim 0.46 M_\odot, \quad D_{OL} \sim 3.3 \text{ kpc}$$


OGLE-2004-BLG-235  
MOA-2004-BLG-53  
(Bond et al 2004)



$$M_p \approx 2.5 M_J, \quad r \approx 4.3 \text{ AU}$$


$$M_* \sim 0.42 M_\odot, \quad D_{OL} \sim 0.2 \text{ kpc}$$

# First Four Detections



Two Jovian-mass planets  
Two Neptune-mass planets


OGLE-2005-BLG-390  
(Beaulieu et al 2006)



$$M_p \sim 5.5 M_{BJ}, \quad r \sim 2.6 \text{ AU}$$

$$M_* \sim 0.55 M_\odot, \quad D_{OL} \sim 0.07 \text{ kpc}$$

OGLE-2005-BLG-169  
(Gould et al 2006)



$$M_p \sim 13 M_{BJ}, \quad r \sim 3.5 \text{ AU}$$

$$M_* = 0.5 M_\odot, \quad D_{OL} = 2.7 \text{ kpc}$$

# Cool Neptunes Are Common

Two low-mass detections imply:

**~37% of stars have Neptunes between 1.6-4.3 AU  
(16-69% at 90% confidence)**

**$dN/d\log a \sim 1$  at  $\sim 3$  AU (0.4-1.6 at 90%)**


**$dN/d\log a \sim 0.3$  at  $\sim 0.15$  AU (HARPS)**

Also:

**Cool Neptunes are more common than cool Jupiters**

# Multiple System

(Gaudi, Bennett et al 2008)



- High-magnification Event
  - $\mu$ FUN, OGLE, MOA
- Must include two planets, finite source, orbital motion, and parallax
- Yields full star and planet masses, information on orbital speed of Saturn and inclination!

High-magnification Event, monitored by  $\mu$ FUN, OGLE, MOA

A  $\sim 0.5 M_{\odot}$  late K-dwarf at  $\sim 1.5$  kpc

Finite  
Source

$$\theta_E \cong 1.48 \text{ mas}$$

Parallax

$$\tilde{r}_E \cong 2.76 \text{ AU}$$

AO Imaging



$$D_l \cong 1.49 \pm 0.13 \text{ kpc}$$

$$M = 0.50 \pm 0.05 M_{\odot}$$



# The OGLE-2006-BLG-109L Planetary System

## Planet b:

$$\text{Mass} = 0.71 \pm 0.08 M_{\text{Jup}}$$


$$\text{Semimajor Axis} = 2.3 \pm 0.2 \text{ AU}$$

## Planet c:


$$\text{Mass} = 0.27 \pm 0.03 M_{\text{Jup}} = 0.90 M_{\text{Sat}}$$

$$\text{Semimajor Axis} = 4.6 \pm 0.5 \text{ AU}$$


# Analog of Jupiter/Saturn



# Implications for Frequency of Systems



(Udalski et al. 2005)





(Bond et al. 2004)

# The Most Massive M Dwarf Planet?

(Dong et al 2008)

Dong et al. 2008



$$M = 0.46 \pm 0.04 M_{\odot}$$

$$D_l = 3.3 \pm 0.4 \text{ kpc}$$

$$v_{\text{LSR}} = 103 \pm 14 \text{ km s}^{-1}$$

$$m = 3.5 \pm 0.3 M_{\text{Jup}}$$


$$r_{\perp} = 3.6 \pm 0.2 \text{ AU}$$

$$T_{eq} \sim 50 K$$

# More on the way...

- **Five or six additional planets in 2007.**
  - Lowest mass planet around lowest mass host (Bennett et al, submitted).
  - Jupiter-mass planet.
  - Another multiple planet system.
- **2008 season underway.**
  - Jupiter-mass planet found two weeks ago.
- **Can expect ~half a dozen planets per year.**

# Current High-Magnification Event



Peak at ~UT 20:30 (~1:30 PDT), magnification  $>1400$   
Sensitive to Earth-mass planets near the Einstein ring.

# What's Next?

- **Current setup (alert/follow-up) saturated**
  - Nearly all of the useable bulge monitored
  - Many events cannot be monitored
  - Monitoring one event at a time too inefficient
- **A new strategy**
  - Dispense with alert/follow-up
  - Simultaneously detect and monitor microlensing events

# What is Required?

- Event Rate

- Primary Event Rate


$$L \approx 10^{-2} \lambda_{-I}$$

- Detection Probability

$$b \approx V^0 \theta^b \approx 10^0 \left( \frac{W^{E_{\text{eff}}}}{W^b} \right)_{I \setminus S}$$

- Detections Per Year

$$N \approx n^E \Omega \Phi L b \approx 10 \lambda_{-I} \left( \frac{10^0 \Omega}{\Omega} \right) \left( \frac{10^0 b \Omega}{\Phi} \right) \left( \frac{10^{-2} \lambda_{-I}}{L} \right) \left( \frac{10^0}{b} \right)$$





# What is Required?

## Detecting the Perturbations from Earth-mass Planets

- Sampling rate  $\sim 10$  minutes

$$\dot{\varphi}^{E,b} = 5\mu_{L2} \left( \frac{W^E}{W^b} \right)_{I \sim 5}$$

- Photometric Accuracy  $\sim 1\%$  at  $I \sim 21$ 
  - Signal Magnitude

$$\frac{\Delta F}{F} \approx 1\% \left( \frac{M_p}{M_\odot} \right) \left( \frac{R_*}{R_\odot} \right)^{-2}$$


- Photometric Uncertainty

$$\sigma = 10^0 \left( \frac{5W}{D} \right)_{-I} \left( \frac{150^2}{\dot{\varphi}^{exb}} \right)_{-I \sim 5} 10^{0.5(1-5I)}$$

# NextGen $\mu$ Lensing Survey

- Requirements to detect  $\sim 10$  Earth-mass planets per year:
  - Monitor  $\sim 10$  square degrees of the Galactic bulge continuously with  $\sim 10$  minute sampling using 1-2m class telescopes, distributed longitudinally throughout the southern hemisphere.
  - Large FOV (2-4 square degree) cameras needed.

# Expected Results



*A next-generation ground-based  $\mu$ lensing survey can test planet formation by probing planets with  $M > M_{\oplus}$  beyond the snow-line.*

# Spontaneous Generation


- MOA-II (NZ, currently operating)
  - 1.8m telescope, 2.18 sq. degree camera
- OGLE -IV (Chile, 2010)
  - 1.3m telescope, upgrade to 1.4 sq. degree camera
- All that is needed is a 1-2m telescope with a large FOV in South Africa.

*“Recommendation A. II. 1 Increase dramatically the efficiency of a ground-based microlensing network by adding a single 2 meter telescope.”*

# Why Space is Better

## From the ground:

- MS sources severely blended
- Getting constraints on hosts is expensive
- Perturbations can be poorly sampled




# What can we expect from Space?

Example: Microlensing Planet Finder (Bennett PI)

- Simulations from Bennett & Rhie (2002)
- Basic results confirmed by independent simulations
- Continuous observations of  $4 \times 0.66$  sq. deg. central Galactic bulge fields:  $\sim 2 \times 10^8$  stars
- Observations in near IR to increase sensitivity
- $\sim 15,000$  events in 4 seasons


# Simulated Planetary Light Curves

- Exposures every 10-15 minutes
- Strong signals
- Unambiguous information
- Moons detectable!  
(1.6 lunar masses)




Bennett & Rhie (2002)

# Wide Range of Separations






# Habitable Planets




- Dozens of Earth-mass habitable planets
- Complements Kepler.

# Free Floating Planets



Planet formation theories generically predict many free-floating planets.


# Lens Detection Provides Accurate Mass Estimate



- Lens will be detected for the majority of main-sequence lenses.
- Host star masses will be measured to 10% for half of the events.
- Projected separations will be measured to 5% for half of the events.

# Planet Detection Sensitivity

- Sensitivity to all Solar System planet analogs except Mercury.
- Demographics of planetary systems - tests planet formation theories.
- Most sensitive technique for  $a \geq 0.5$  AU.
- Good sensitivity to “outer” habitable zone (Mars-like orbits) where detection by imaging is easiest.
- Complementary to Kepler.
- Assumes  $\geq 9\sigma$  detection threshold.
- Can find moons and free floating planets.



Updated from Bennett & Rhie (2002) ApJ 574, 985

*“Recommendation B. II. 2 Without impacting the launch schedule of the astrometric mission cited above\*, launch a Discovery-class space-based microlensing mission to determine the statistics of planetary mass and the separation of planets from their host stars as a function of stellar type and location in the galaxy, and to derive  $\eta_{\oplus}$  over a very large sample.*

*\* “Recommendation B. I. a. 1 Launch and operate a space based astrometric mission capable of detecting planets down to the mass of the Earth around 60-100 nearby stars...”*

# Technology

- Ground-based 1-2m, Wide FOV Telescope
  - Several very similar telescopes already operating
    - MOA-II
    - Pan-STARRS-1 - \$20M
- Space-based microlensing mission
  - Requires almost no technology development.
  - Can extensively leverage other missions (Spitzer, NextView, Ikonos, JWST)
  - Can use many components that are demonstrated on orbit or flight qualified.

# MPF Mission Design

- 1.1-m aperture consisting of a three-mirror anastigmat telescope feeding a 147 Mpixel HgCdTe focal plane (35 2048<sup>2</sup> arrays)
- The spacecraft bus is a near-identical copy of that used for *Spitzer*.
- The telescope system very similar to NextView commercial Earth-observing telescope designs.
- Detectors developed for JWST meet MPFs requirements.
- All elements are at TRL 6 or better.
- Total Cost \$300 (without launch vehicle)

Property	Value	Units
Launch Vehicle	7920-9.5	Delta II
Orbit	Inclined GEO 28.7	degrees
Mission Lifetime	4.0	years
Telescope Aperture	1.1	meters (diam.)
Field of View	0.95x0.68	degrees
Spatial Resolution	0.240	arcsec/pixel
Pointing Stability	0.048	arcsec
Focal Plane Format	145	Megapixels
Spectral Range	600-1700	nm in 3 bands
Quantum Efficiency	>75% >55%	900-1400 nm 700-1600 nm
Dark Current	<1	e-/pixel/sec
Readout Noise	<30	e-/read
Photometric Accuracy	1% or better	at J=20.5
Data Rate	50.1	Mbits/sec
MPF Mission Requirements		

# Dark Energy Synergy

- Space-based microlensing mission telescope requirements are very similar to the requirements for many proposed dark energy missions.
- Combined dark energy/planet finding mission probably could be accomplished at a substantial savings.
- ADEPT, Destiny, SNAP, DUNE/SPACE/Euclid
  - Wide FOV, >1.1m aperture, technical specifications appear to satisfy space-based microlensing survey specifications
  - DUNE/SPACE/Euclid can meet all the science goals without modification to hardware.
- Trade study:
  - Observing time
  - Pass bands
  - FOV and Detectors
  - Orbit
  - Telemetry
  - Aperture
  - Optics
  - Pointing



# Summary

- Ground-based Next-Generation Survey:     +\$10M—\$20M
  - Complete network with a single wide FOV 1-2m telescope in SA.
  - Frequency of planets  $>M_{\oplus}$  beyond the snow line.
  - Test planet formation theories.
- **Either:** Space-based Microlensing Mission: +\$300M + launch
  - Complete census of planets with mass greater than Mars and  $a > 0.5$  AU.
  - Sensitivity to all Solar System planet analogs except Mercury.
  - Demographics of planetary systems - tests planet formation theories.
  - Detect “outer” habitable zone (Mars-like orbits) where detection by imaging is easiest.
  - Can find moons and free floating planets.
- **Or:** Joint  $\mu$ lensing/Dark Energy Mission     +\$100M—\$200M?
- Total cost to “Exoplanet Community”: **\$120M—\$420M**